arXiv:hep-ph/9807364v2 8 Sep 1998

New CP-violating parameters in cascade decays

A. Amorim[a,b], Mário G. Santos[a] and João P. Silva[a,c]

[a] Centro de Física Nuclear da Universidade de Lisboa
Avenida Professor Gama Pinto, 2,
1699 Lisboa Codex, Portugal

[b] Departamento de Física
Faculdade de Ciências da Universidade de Lisboa
Campo Grande, 1700 Lisboa, Portugal

[c] Centro de Física
Instituto Superior de Engenharia de Lisboa
1900 Lisboa, Portugal

February 1, 2008

Abstract

We consider decay chains of the type $P \to M + \cdots \to f + \cdots$, where M is a neutral meson that may mix with its antiparticle $\overline{M}$, before decaying into the final state f. P may be either a heavier neutral meson or a charged meson. We perform a rephasing-invariant analysis of the quantities that show up in such cascade decays. If the decay $P \to M + \cdots$ (or the decay $P \to \overline{M} + \cdots$) is forbidden, we find the usual λ_f parameters describing the interference between the mixing of a neutral meson system and the decay *from that system* into the final state f. However, when both the $P \to M + \cdots$ and $P \to \overline{M} + \cdots$ decays are allowed, we find a new class of rephasing-invariant parameters, ξ_i, that measure the interference between the mixing of a neutral meson system and the decay from the initial state *into that system*. We show that the quantities λ_f and ξ_i are necessary and sufficient to describe all the interference effects present in the most general cascade decay. We discuss the various cascade decays in turn, highlighting the special features of each one.

1 Introduction

The particle-antiparticle neutral meson systems have long been identified as an ideal setting to search for CP violation (CPV). This effect was discovered in 1964 in the neutral kaon system [1], and it is believed that this will soon be complemented with measurements of CPV in the decays of neutral B mesons.

The CP violation found in the kaon system and the null results found elsewhere can be accommodated in the Standard Model (SM), through an irremovable complex phase in the Cabibbo-Kobayashi-Maskawa quark mixing matrix [2]. The major goal of the upcoming B-factory experiments is to submit this explanation to stringent tests [3].

In 1989, Azimov [4] pointed out that additional tests could be performed by looking for $B_d \to J/\psi K \to J/\psi [f]_K$ 'cascade decays', involving the neutral B_d and neutral kaon systems in succession. This idea has been followed by Dass and Sarma [5]. Recently, cascade decays have received renewed attention. Kayser and Stodolsky have developed a matrix method to deal with cascade decays [6]; Kayser has shown that one may use $B_d \to J/\psi K \to J/\psi [\pi l \nu]_K$ decays to get at $\cos 2\beta$ [7]; and Azimov and Dunietz pointed out that one may use $B_s \to J/\psi K \to J/\psi [f]_K$ decays to measure Δm_{B_s}, even if Δm_{B_s} turns out to be too large to be determined experimentally from the direct decays of the B_s mesons [8].

In all these cases, an initial B^0 meson[1] can only decay to one of the kaon's flavour eigenstates. To leading order in the SM, the decays $B_d^0 \to \overline{K^0} + X$ and $B_s^0 \to K^0 + X$, and the respective CP conjugate decays are forbidden. In this article we will consider a more general situation, in which a given meson can decay into both flavour eigenstates of a lighter neutral meson system. This situation is the one relevant for the decays

$$\begin{aligned} B^\pm &\to D + X^\pm \to [f]_D + X^\pm, \qquad & B &\to D + X \to [f]_D + X, \\ D^\pm &\to K + X^\pm \to [f]_K + X^\pm, \qquad & D &\to K + X \to [f]_K + X. \end{aligned} \tag{1}$$

Decays of the first type have been discussed by Meca and Silva [9] in the context of uncovering new physics effects in the $D^0 - \overline{D^0}$ system. Here we show that all these decays depend on a new type of CP-violating observables, over and above those required for the description of direct decays of neutral mesons.

In section 2 we describe all the quantities involved in cascade decays of the type $P \to M + X \to [f]_M + X$, where P and M describe two neutral meson systems, and $[f]_M$ describes a set of particles into which M can decay, and which have an invariant mass equal to that of the $M^0 - \overline{M^0}$ system. This analysis is performed by noting that all physical quantities must remain invariant under a rephasing of the kets in the problem. In section 3 we develop the conditions for CP invariance, thus identifying all the CP-violating rephasing-invariant quantities involved in cascade decays. Naturally, we obtain the usual CP-violating observables. These describe CPV in the mixing of the neutral meson systems; CPV in the decays; and CPV in the interference between the mixing of the neutral meson system and the decay *from that system* into the final state. However, when both the $P \to M + \cdots$ and $P \to \overline{M} + \cdots$ decays are allowed, we find a new class of rephasing invariant quantities that measure the interference between the mixing of the intermediate neutral meson system M, and the decay from the initial state P *into that system*. These quantities are used in section 4 to develop the formulae describing a generic cascade decay. Section 5 contains a classification of cascade decays and an extended discussion of the special features of each case. We draw our conclusions in section 6.

---

[1] We will take B^0 to stand for both B_d^0 and B_s^0. Moreover, K refers to a generic combination of K^0 and $\overline{K^0}$, and similarly for D, B_d and B_s.

2 Rephasing-invariant quantities

Let us consider a decay of the type

$$P \to M + X \to [f]_M + X, \tag{2}$$

where both P and M belong to neutral meson systems. In general, the amplitude for the overall decay chain involves the amplitudes of the primary decays

$$\begin{aligned} A_{P^0\to M^0} &\equiv \langle M^0 X|T|P^0\rangle\ , & A_{\overline{P^0}\to M^0} &\equiv \langle M^0 X|T|\overline{P^0}\rangle\ , \\ A_{P^0\to \overline{M^0}} &\equiv \langle \overline{M^0} X|T|P^0\rangle\ , & A_{\overline{P^0}\to \overline{M^0}} &\equiv \langle \overline{M^0} X|T|\overline{P^0}\rangle\ , \end{aligned} \tag{3}$$

followed by the amplitudes of the secondary decay,

$$A_{M^0\to f} \equiv \langle f|T|M^0\rangle\ , \quad A_{\overline{M^0}\to f} \equiv \langle f|T|\overline{M^0}\rangle\ . \tag{4}$$

Since there is mixing in the neutral $P^0 - \overline{P}^0$ system, the decay rate will also involve the parameter q_P/p_P which arises in the transformation from the flavour eigenstates into the mass eigenstates:

$$\begin{aligned} |P_H\rangle &= p_P|P^0\rangle + q_P|\overline{P^0}\rangle, \\ |P_L\rangle &= p_P|P^0\rangle - q_P|\overline{P^0}\rangle. \end{aligned} \tag{5}$$

Here, P_H and P_L refer to the heavy and light mass eigenstates, respectively. Similarly, the $M^0 - \overline{M^0}$ mixing enters in the expressions through the parameter q_M/p_M. For completeness, we describe in appendix A the time evolution of a generic neutral meson system.

In quantum mechanics, all kets may be rephased at will,[2]

$$\begin{aligned} |P^0\rangle &\to e^{i\gamma_P}|P^0\rangle\ , & |\overline{P^0}\rangle &\to e^{i\overline{\gamma}_P}|\overline{P^0}\rangle\ , \\ |M^0\rangle &\to e^{i\gamma_M}|M^0\rangle\ , & |\overline{M^0}\rangle &\to e^{i\overline{\gamma}_M}|\overline{M^0}\rangle\ , \\ |f\rangle &\to e^{i\gamma_f}|f\rangle\ , & |\bar{f}\rangle &\to e^{i\overline{\gamma}_f}|\bar{f}\rangle\ . \end{aligned} \tag{6}$$

Under this rephasing, the amplitudes and mixing parameters also change, according to

$$\begin{aligned} \tfrac{q_P}{p_P} &\to e^{i(\gamma_P-\overline{\gamma}_P)}\tfrac{q_P}{p_P}\ , & \tfrac{q_M}{p_M} &\to e^{i(\gamma_M-\overline{\gamma}_M)}\tfrac{q_M}{p_M}\ , \\ A_{P^0\to M^0} &\to e^{i(\gamma_P-\gamma_M)}A_{P^0\to M^0}\ , & A_{\overline{P^0}\to M^0} &\to e^{i(\overline{\gamma}_P-\gamma_M)}A_{\overline{P^0}\to M^0}\ , \\ A_{P^0\to \overline{M^0}} &\to e^{i(\gamma_P-\overline{\gamma}_M)}A_{P^0\to \overline{M^0}}\ , & A_{\overline{P^0}\to \overline{M^0}} &\to e^{i(\overline{\gamma}_P-\overline{\gamma}_M)}A_{\overline{P^0}\to \overline{M^0}}\ , \\ A_{M^0\to f} &\to e^{i(\gamma_M-\gamma_f)}A_{M^0\to f}\ , & A_{\overline{M^0}\to f} &\to e^{i(\overline{\gamma}_M-\gamma_f)}A_{\overline{M^0}\to f}\ , \\ A_{M^0\to \bar{f}} &\to e^{i(\gamma_M-\overline{\gamma}_f)}A_{M^0\to \bar{f}}\ , & A_{\overline{M^0}\to \bar{f}} &\to e^{i(\overline{\gamma}_M-\overline{\gamma}_f)}A_{\overline{M^0}\to \bar{f}}\ . \end{aligned} \tag{7}$$

---

[2] The rephasing-invariant analysis of cascade decays presented in this section is inspired in the analysis of direct decays performed in Ref. [10].

Only those quantities which are invariant under this redefinition may have physical meaning. Clearly, the magnitudes of all the quantities in Eq. 7 satisfy this condition.

In addition, we find some rephasing-invariant quantities which arise from the clash between the phases in the mixing and the phases in the decay amplitudes:

$$\lambda_{P\to M^0} \equiv \frac{q_P}{p_P}\frac{A_{\overline{P^0}\to M^0}}{A_{P^0\to M^0}}\ , \qquad \lambda_{P\to\overline{M^0}} \equiv \frac{q_P}{p_P}\frac{A_{\overline{P^0}\to\overline{M^0}}}{A_{P^0\to\overline{M^0}}}\ , \tag{8}$$

$$\lambda_{M\to f} \equiv \frac{q_M}{p_M}\frac{A_{\overline{M^0}\to f}}{A_{M^0\to f}}\ , \qquad \lambda_{M\to\bar f} \equiv \frac{q_M}{p_M}\frac{A_{\overline{M^0}\to\bar f}}{A_{M^0\to\bar f}}\ , \tag{9}$$

$$\xi_{P^0\to M} \equiv \frac{A_{P^0\to\overline{M^0}}}{A_{P^0\to M^0}}\frac{p_M}{q_M}\ , \qquad \xi_{\overline{P^0}\to M} \equiv \frac{A_{\overline{P^0}\to\overline{M^0}}}{A_{\overline{P^0}\to M^0}}\frac{p_M}{q_M}\ . \tag{10}$$

The quantities in Eq. 8 are well known. They describe the interference between the mixing in the $P^0-\overline{P^0}$ system and the decay *from that system* into the flavour specific meson M^0 or $\overline{M^0}$, respectively. Similarly, the parameters in Eq. 9 describe the interference between the mixing in the $M^0-\overline{M^0}$ system and the decay *from that system* into the final state f and $\bar f$, respectively. On the contrary, the parameters in Eq. 10 describe a new type of interference: the interference between the mixing in the $M^0-\overline{M^0}$ system and the decay *into that system* (originating from P^0, in the case of $\xi_{P^0\to M}$, or from $\overline{P^0}$, in the case of $\xi_{\overline{P^0}\to M}$).

We notice that the parameters described thus far are not all independent. In particular

$$\lambda_{P\to M^0}\xi_{\overline{P^0}\to M} = \lambda_{P\to\overline{M^0}}\xi_{P^0\to M}. \tag{11}$$

This means that, of the six phases contained in Eqs. 8, 9, and 10, only five are independent. This is obvious from Eqs. 7 which contain ten quantities and five rephasings.

3 Conditions for CP-invariance

In order to study CP violation we first consider the consequences of CP invariance. If CP is a good symmetry, then there exist three phases α_P, α_M and α_f, and two CP eigenvalues $\eta_P = \pm 1$ and $\eta_M = \pm 1$, such that:

$$\begin{aligned}\mathcal{CP}|P^0\rangle &= e^{i\alpha_P}|\overline{P^0}\rangle\ ,\\ \mathcal{CP}|M^0\rangle &= e^{i\alpha_M}|\overline{M^0}\rangle\ ,\\ \mathcal{CP}|f\rangle &= e^{i\alpha_f}|\bar f\rangle\ ,\end{aligned} \tag{12}$$

and

$$\begin{aligned}\mathcal{CP}|P_H\rangle &= \eta_P|P_H\rangle\ , & \mathcal{CP}|P_L\rangle &= -\eta_P|P_L\rangle\ ,\\ \mathcal{CP}|M_H\rangle &= \eta_M|M_H\rangle\ , & \mathcal{CP}|M_L\rangle &= -\eta_M|M_L\rangle\ .\end{aligned} \tag{13}$$

Notice that, since we define the sign of Δm to be positive, it is the value of η_P and of η_M that must be taken from experiment. For example, we know experimentally that, if the small CP violation in the neutral kaon system were absent, then the heavy kaon would coincide with the long-lived kaon, which would be CP odd. Therefore, we must take $\eta_K = -1$ whenever we neglect CP violation in the neutral kaon system. For the other neutral meson systems, these parameters must still be determined experimentally. In the meantime, one sometimes uses the SM predictions for these parameters. For instance, the SM prediction for the $B_s^0 - \overline{B_s^0}$ system is that the heavy state should be mostly CP odd [11]. In this case, neglecting the CP violation in the $B_s^0 - \overline{B_s^0}$ system leads to $\eta_{B_s} = -1$.

Moreover, the CP transformation of the composite intermediate state is

$$\mathcal{CP}|XM^0\rangle = \eta_X e^{i\alpha_M} |\overline{X}\,\overline{M^0}\rangle \ . \tag{14}$$

Here η_X contains any relevant CP transformation of the state X, as well as the parity corresponding to the relative angular momentum between X and M. For example, in the decays $B \to J/\psi K$, $\eta_X = -1$ because J/ψ is CP even, while the J/ψ and the kaon have a relative $L = 1$ orbital angular momentum [12].

As a result of Eqs. 12, 13 and 14, the CP invariance conditions are

$$\frac{q_P}{p_P} = \eta_P e^{i\alpha_P} \ , \qquad \frac{q_M}{p_M} = \eta_M e^{i\alpha_M} \ , \tag{15}$$

and

$$\begin{array}{rclrcl} A_{P^0\to M^0} & = & \eta_X e^{i(\alpha_P - \alpha_M)} A_{\overline{P^0}\to\overline{M^0}} \ , & A_{P^0\to\overline{M^0}} & = & \eta_X e^{i(\alpha_P + \alpha_M)} A_{\overline{P^0}\to M^0} \ , \\ A_{M^0\to f} & = & e^{i(\alpha_M - \alpha_f)} A_{\overline{M^0}\to\bar{f}} \ , & A_{M^0\to\bar{f}} & = & e^{i(\alpha_M + \alpha_f)} A_{\overline{M^0}\to f} \ . \end{array} \tag{16}$$

Therefore, if CP is conserved we have

$$\begin{array}{rclrcl} \left|\frac{q_P}{p_P}\right| & = & 1 & \left|\frac{q_M}{p_M}\right| & = & 1 \\ |A_{P^0\to M^0}| & = & \left|A_{\overline{P^0}\to\overline{M^0}}\right| & \left|A_{P^0\to\overline{M^0}}\right| & = & \left|A_{\overline{P^0}\to M^0}\right| \\ |A_{M^0\to f}| & = & \left|A_{\overline{M^0}\to\bar{f}}\right| & \left|A_{M^0\to\bar{f}}\right| & = & \left|A_{\overline{M^0}\to f}\right| \ . \end{array} \tag{17}$$

Moreover, the parameters describing interference CP violation become related by

$$\begin{array}{rcl} \lambda_{P\to M^0}\,\lambda_{P\to\overline{M^0}} & = & 1 \\ \lambda_{M\to f}\,\lambda_{M\to\bar{f}} & = & 1 \\ \xi_{P^0\to M}\,\xi_{\overline{P^0}\to M} & = & 1 \\ \lambda_{P\to M^0}\,\xi_{\overline{P^0}\to M} = \lambda_{P\to\overline{M^0}}\,\xi_{P^0\to M} & = & \eta_X\eta_P\eta_M . \end{array} \tag{18}$$

We have used Eq. 11 on the last line. One may use the relations above to develop more complicated conditions for CP invariance, such as

$$A_{\overline{P^0}\to\overline{M^0}} A_{\overline{M^0}\to f} A^*_{\overline{P^0}\to M^0} A^*_{M^0\to f} = A_{P^0\to M^0} A_{M^0\to\bar{f}} A^*_{P^0\to\overline{M^0}} A^*_{\overline{M^0}\to\bar{f}} . \tag{19}$$

A very important special case occurs when f is a CP eigenstate. Then $\eta_f \equiv e^{i\alpha_f} = \pm 1$, and the conditions for CP invariance become

$$|A_{M^0\to f}| = \left|A_{\overline{M^0}\to f}\right| \quad \text{and} \quad \lambda_{M\to f} = \eta_M\eta_f . \tag{20}$$



Figure 1: The most general $P \to MX \to [f]_M X$ cascade decay.

We have found the usual types of CP violation: $|q/p| - 1$ probes CP violation in the mixing of neutral mesons; $|A_{i\to f}| - |A_{\bar{i}\to\bar{f}}|$ probes CP violation in the decays; while $\arg\lambda_{i\to f} + \arg\lambda_{i\to\bar{f}}$ measures CP violation arising in the interference between the mixing of the initial neutral meson system, i, and the decay *from that system* into the final states f and $\bar{f}$. If f is a CP eigenstate, then the CP-violating observable $\arg\lambda_{i\to f} + \arg\lambda_{i\to\bar{f}}$ is related to $\mathrm{Im}\lambda_{i\to f}$. In addition, we have discovered a new type of CP-violating observables given by $\arg\xi_{P^0\to M} + \arg\xi_{\overline{P^0}\to M}$. These observables measure the CP violation which arises from the interference between the mixing in the $M^0 - \overline{M^0}$ system and the decay from the initial state (P^0 and $\overline{P^0}$) *into that system.*

4 The decay rate for a generic cascade decay

4.1 Decay amplitudes

Let us consider a neutral meson P^0 that decays into an intermediate state containing a combination of M^0 and $\overline{M^0}$ at time t_P, measured in the rest-frame of P^0. The resulting combination of M^0 and $\overline{M^0}$ will evolve in time and decay into the final state f at time t_M, measured in its rest-frame. The decay amplitude for this process is given by

$$\begin{aligned}
A\left[P^0 \overset{t_P}{\to} M \overset{t_M}{\to} f\right] &= g_+^P(t_P) g_+^M(t_M) \left[A_{P^0\to M^0} A_{M^0\to f} + A_{P^0\to\overline{M^0}} A_{\overline{M^0}\to f}\right] \\
&+ g_+^P(t_P) g_-^M(t_M) \left[\frac{q_M}{p_M} A_{P^0\to M^0} A_{\overline{M^0}\to f} + \frac{p_M}{q_M} A_{P^0\to\overline{M^0}} A_{M^0\to f}\right] \\
&+ g_-^P(t_P) g_-^M(t_M) \frac{q_P}{p_P} \left[\frac{p_M}{q_M} A_{\overline{P^0}\to\overline{M^0}} A_{M^0\to f} + \frac{q_M}{p_M} A_{\overline{P^0}\to M^0} A_{\overline{M^0}\to f}\right] \\
&+ g_-^P(t_P) g_+^M(t_M) \frac{q_P}{p_P} \left[A_{\overline{P^0}\to\overline{M^0}} A_{\overline{M^0}\to f} + A_{\overline{P^0}\to M^0} A_{M^0\to f}\right]. \qquad (21)
\end{aligned}$$

This result is easiest to derive with the help of the evolution diagram presented in Fig. 1. Henceforth we shall not show the explicit time dependence of the g-functions.

The decay amplitude may also be written as

$$\begin{aligned}
A\left[P^0 \overset{t_P}{\to} M \overset{t_M}{\to} f\right] &= \left(A_{P^0\to M^0} A_{M^0\to f} + A_{P^0\to\overline{M^0}} A_{\overline{M^0}\to f}\right) \times \\
&\quad \left[g_+^P g_+^M + \chi_1\, g_+^P g_-^M + \chi_2\, g_-^P g_-^M + \chi_3 g_-^P g_+^M\right] \ , \qquad (22)
\end{aligned}$$

where we have used

$$\begin{aligned}
\chi_1 &\equiv \frac{A_{P^0\to M^0}\frac{q_M}{p_M}A_{\overline{M^0}\to f}+A_{P^0\to\overline{M^0}}\frac{p_M}{q_M}A_{M^0\to f}}{A_{P^0\to M^0}A_{M^0\to f}+A_{P^0\to\overline{M^0}}A_{\overline{M^0}\to f}} &&= \frac{\lambda_{M\to f}+\xi_{P^0\to M}}{1+\xi_{P^0\to M}\lambda_{M\to f}}\ , \\
\chi_2 &\equiv \frac{q_P}{p_P}\frac{A_{\overline{P^0}\to\overline{M^0}}\frac{p_M}{q_M}A_{M^0\to f}+A_{\overline{P^0}\to M^0}\frac{q_M}{p_M}A_{\overline{M^0}\to f}}{A_{P^0\to M^0}A_{M^0\to f}+A_{P^0\to\overline{M^0}}A_{\overline{M^0}\to f}} &&= \lambda_{P\to M^0}\frac{\xi_{\overline{P^0}\to M}+\lambda_{M\to f}}{1+\xi_{P^0\to M}\lambda_{M\to f}}\ , \\
\chi_3 &\equiv \frac{q_P}{p_P}\frac{A_{\overline{P^0}\to\overline{M^0}}A_{\overline{M^0}\to f}+A_{\overline{P^0}\to M^0}A_{M^0\to f}}{A_{P^0\to M^0}A_{M^0\to f}+A_{P^0\to\overline{M^0}}A_{\overline{M^0}\to f}} &&= \lambda_{P\to M^0}\frac{\xi_{\overline{P^0}\to M}\lambda_{M\to f}+1}{1+\xi_{P^0\to M}\lambda_{M\to f}}\ .
\end{aligned} \tag{23}$$

Notice that the parameters χ_n involve $\xi_{P^0\to M}$ and $\xi_{\overline{P^0}\to M}$, which describe the new type of interference effects which are the subject of this article.

However, χ_3 may always be reinterpreted in a different way. Let us consider the state

$$|M_{\text{into}\,f}\rangle \equiv A^*_{M^0\to f}|M^0\rangle + A^*_{\overline{M^0}\to f}|\overline{M^0}\rangle\ . \tag{24}$$

This is the $M^0-\overline{M^0}$ combination that decays into the final state f at time t_M. Indeed, this state is orthogonal to the state

$$|M_{\text{not into}\,f}\rangle \equiv A_{\overline{M^0}\to f}|M^0\rangle - A_{M^0\to f}|\overline{M^0}\rangle\ , \tag{25}$$

which clearly cannot decay into the final state f because $\langle f|T|M_{\text{not into}\,f}\rangle = 0$. Using Eq. 24, it is easy to show that

$$\chi_3 = \lambda_{P\to M_{\text{into}\,f}}. \tag{26}$$

Still, in general, χ_1 and χ_2 may not be reinterpreted as parameters λ.

One may wonder why these effects did not show up in the analysis presented in [4, 5, 6, 7, 8]. The reason is simple: in those cases a meson P^0 can only decay into one of the flavour eigenstates of the intermediate meson system. For example, P^0 can decay into M^0, but not into $\overline{M^0}$. Indeed, taking $A_{\overline{P^0}\to M^0} = 0 = A_{P^0\to\overline{M^0}}$, we find, in addition to Eq. 26,

$$\chi_1 = \lambda_{M\to f} \quad \text{and} \quad \chi_2 = \lambda_{P\to M_H}. \tag{27}$$

The last equality involves M_H, the heavy mass eigenstate of the $M^0-\overline{M^0}$ system, and holds if and only if $|p_M/q_M| = 1$. In these cases, we do not need to introduce the parameters ξ. On the contrary, these parameters are required in the discussion of the $B^+\to DK^+\to[f]_D K^+$ cascade decays, as found by Meca and Silva [9]. The same parameters would enter the analysis of the decays, $B^+\to D_{H,L}K^+$, if only these decays could be disentangled from one another [13]. Unfortunately, they cannot [13]. It should now be clear that the effects described by $\xi_{P^0\to M}$ and $\xi_{\overline{P^0}\to M}$ are mandatory and, when combined with the usual CP-violating observables, are also sufficient for a discussion of the $B\to D$ and $D\to K$ decay chains.

In the example just discussed, in which $A_{\overline{P^0}\to M^0} = 0 = A_{P^0\to\overline{M^0}}$, we find that $\chi_3 = \chi_1\chi_2$. This relation, although not valid in general, holds in a variety of special cases. Indeed,

$$\chi_3 - \chi_1\chi_2 = \frac{\left(1-\lambda^2_{M\to f}\right)\left(1-\xi_{\overline{P^0}\to M}\xi_{P^0\to M}\right)\lambda_{P\to M^0}}{\left(1+\xi_{P^0\to M}\lambda_{M\to f}\right)^2}. \tag{28}$$

This result is completely general. We conclude that $\chi_3-\chi_1\chi_2$ vanishes if any of the following conditions holds:

1. $\lambda^2_{M\to f} = 1$, which, if f is a CP eigenstate, means that there is CP conservation when the $M^0 - \overline{M^0}$ system decays into the final state f;

2. $\xi_{\overline{P^0}\to M}\xi_{P^0\to M} = 1$, meaning that there is CP conservation when the $P^0 - \overline{P^0}$ system decays into the $M^0 - \overline{M^0}$ system;

3. $\lambda_{P\to M^0} = 0$, meaning that the decay $\overline{P^0} \to M^0 X$ is forbidden. Analogously, the difference also vanishes if $P^0 \to M^0 X$ is forbidden.

If any of these conditions holds, the amplitude for the cascade decay depends only on two independent parameters (for example χ_1 and χ_2), in addition to the overall normalization factor. We see that $\chi_3 - \chi_1\chi_2$ is only different from zero if there are interference effects in every step of the decay chain.[3]

We now turn to the CP invariance constraints on the χ_n parameters. We find that, if there is CP conservation, then

$$\chi_2 = \eta_X \eta_P \eta_M , \tag{29}$$

and $\chi_3 = \chi_1\chi_2$. In addition, if f is a CP eigenstate, then we find two further conditions for CP invariance:

$$\chi_1 = \eta_M \eta_f \quad \text{and} \quad \chi_3 = \eta_X \eta_P \eta_f . \tag{30}$$

We now consider the case in which we start with a tagged $\overline{P^0}$. The decay amplitude for the cascade chain becomes

$$\begin{aligned} A\left[\overline{P^0} \overset{t_P}{\to} M \overset{t_M}{\to} f\right] &= \frac{p_P}{q_P} \left(A_{P^0\to M^0} A_{M^0\to f} + A_{P^0\to\overline{M^0}} A_{\overline{M^0}\to f}\right) \times \\ & \left[\chi_3\, g^P_+ g^M_+ + \chi_2\, g^P_+ g^M_- + \chi_1\, g^P_- g^M_- + g^P_- g^M_+\right] . \end{aligned} \tag{31}$$

This result may be obtained from Eq. 22 by interchanging $g^P_+ \leftrightarrow g^P_-$ and multiplying the result by p_P/q_P.

4.2 Decay rates

Using Eq. 22, we find for the decay $P^0 \to MX \to [f]_M X$,

$$\begin{aligned} \Gamma\left[P^0 \overset{t_P}{\to} M \overset{t_M}{\to} f\right] &= N \Big[\left|g^P_+ g^M_+\right|^2 + |\chi_1|^2 \left|g^P_+ g^M_-\right|^2 + |\chi_2|^2 \left|g^P_- g^M_-\right|^2 + |\chi_3|^2 \left|g^P_- g^M_+\right|^2 \\ &+ 2\left|g^P_+\right|^2 \text{Re}\left\{\chi_1 g^{M*}_+ g^M_-\right\} + 2\left|g^P_-\right|^2 \text{Re}\left\{\chi_2\chi_3^* g^{M*}_+ g^M_-\right\} \\ &+ 2\left|g^M_+\right|^2 \text{Re}\left\{\chi_3 g^{P*}_+ g^P_-\right\} + 2\left|g^M_-\right|^2 \text{Re}\left\{\chi_1^*\chi_2 g^{P*}_+ g^P_-\right\} \\ &+ 2\text{Re}\left\{\chi_2 g^{P*}_+ g^P_- g^{M*}_+ g^M_-\right\} + 2\text{Re}\left\{\chi_1^*\chi_3 g^{P*}_+ g^P_- g^M_+ g^{M*}_-\right\}\Big] , \end{aligned} \tag{32}$$

where

$$N \equiv \left|A_{P^0\to M^0} A_{M^0\to f} + A_{P^0\to\overline{M^0}} A_{\overline{M^0}\to f}\right|^2 . \tag{33}$$

---

[3] Informally, we may think of $\chi_3 - \chi_1\chi_2$ as the “all hell breaks loose” parameter.



Figure 2: The $B^+ \to DX^+ \to (K^-\pi^+)_D X^+$ cascade decay.

Similarly, the decay rate for an initial $\overline{P^0}$ is

$$\begin{aligned}
\frac{\Gamma\left[\overline{P^0} \overset{t_P}{\to} M \overset{t_M}{\to} f\right]}{|p_P/q_P|^2} &= N\Big[|\chi_3|^2 \left|g_+^P g_+^M\right|^2 + |\chi_2|^2 \left|g_+^P g_-^M\right|^2 + |\chi_1|^2 \left|g_-^P g_-^M\right|^2 + \left|g_-^P g_+^M\right|^2 \\
&+ 2\left|g_+^P\right|^2 \mathrm{Re}\,\left\{\chi_2\chi_3^* g_+^{M*} g_-^M\right\} + 2\left|g_-^P\right|^2 \mathrm{Re}\,\left\{\chi_1 g_+^{M*} g_-^M\right\} \\
&+ 2\left|g_+^M\right|^2 \mathrm{Re}\,\left\{\chi_3^* g_+^{P*} g_-^P\right\} + 2\left|g_-^M\right|^2 \mathrm{Re}\,\left\{\chi_1\chi_2^* g_+^{P*} g_-^P\right\} \\
&+ 2\mathrm{Re}\,\left\{\chi_1\chi_3^* g_+^{P*} g_-^P g_+^{M*} g_-^M\right\} + 2\mathrm{Re}\,\left\{\chi_2^* g_+^{P*} g_-^P g_+^M g_-^{M*}\right\}\Big]\ , \qquad (34)
\end{aligned}$$

One may also use the rates in Eqs. 32 and 34 to describe decays of the type $P^\pm \to MX^\pm \to [f]_M X^\pm$. To this end, one substitutes $P^0 \to P^+$, $\overline{P^0} \to P^-$, $g_+^P \to \exp\left(-\Gamma_{P^+} t_P/2\right)$, $g_-^P \to 0$, and $q_P/p_P \to 1$.

5 Classification of cascade decays

To simplify the notation we will suppress the explicit reference to the final stage of the decay chain, $[f]_M X$, unless it is required. Traditionally, one has considered cascade decays of the type $B_d, B_s \to K$, using the known mixing parameters of the $K^0 - \overline{K^0}$ system as an analyzer for the parameters of the heavier $B_d^0 - \overline{B_d^0}$ [4, 5, 6, 7] and $B_s^0 - \overline{B_s^0}$ [8] systems. More recently, Meca and Silva [9] have stressed the role of $B^\pm \to D$ decays as a probe for new physics contributions to the $D^0 - \overline{D^0}$ mixing.

Clearly, cascade decays of the type $P^\pm \to MX^\pm \to [f]_M X^\pm$ are useful for the second strategy, while cascade decays of the type $P \to MX \to [f]_M X$ could, in principle, be useful for both. We analyze them in turn.

5.1 Cascade decays from charged P mesons

Decays of this type include $D^+ \to \overline{K^0} K^+$, $D^+ \to \{\overline{K^0}, K^0\}\pi^+$, $D_s^+ \to K^0\pi^+$, and $D_s^+ \to \{K^0, \overline{K^0}\}K^+$. Since the kaon system is well tested experimentally, these decays are of limited use. This is also true of the pure penguin decays $B^+ \to K^0\pi^+$ and $B^+ \to \overline{K^0}K^+$.

More interesting are the decays $B^+ \to \{D^0, \overline{D^0}\}K^+$, such as the one represented in Fig. 2. We use

$$\epsilon \equiv \left|\frac{A_{\overline{D^0}\to K^-\pi^+}}{A_{D^0\to K^-\pi^+}}\right| \sim \left|\frac{V_{cd}V_{us}}{V_{cs}V_{ud}}\right| \sim 0.05 \ , \tag{35}$$

and [14]

$$\tilde{\epsilon} \equiv \left|\frac{A_{B^+\to D^0 K^+}}{A_{B^+\to \overline{D^0} K^+}}\right| \sim \left|\frac{V_{ub}V_{cs}}{V_{cb}V_{us}}\frac{a_2}{a_1}\right| \sim 0.09 \ , \tag{36}$$

where $|a_2/a_1| \sim 0.26$ accounts for the fact that the $B^+ \to D^0 K^+$ decay is color-suppressed, while the $B^+ \to \overline{D^0} K^+$ decay is not.

Meca and Silva have pointed out that these decays may be used to get at $x_D \equiv \Delta m_D/\Gamma_D$ [9]. This is easy to see from Fig. 2, where we have taken $y_D = 0$, and used the small x_D approximation, $g_+^D \sim 1$ and $g_-^D \sim -i/2\, x_D \tau_D$, with $\tau_D \equiv \Gamma_D t_D$. If $\Delta m_D = 0$, then there are no mixed decay paths and we only have two contributions to the decay amplitude. In suitable units, one is of order ϵ and the other of order $\tilde{\epsilon}$. On the other hand, if $\Delta m_D \neq 0$, then there are two further decay paths; one is of order x_D and the other of order $\epsilon\tilde{\epsilon}x_D$. Now, values of $x_D \sim 10^{-2}$ are easy to obtain in many models of new physics [15]. In that case, the last term may be dropped, but the term linear in x_D corrects the unmixed ϵ and $\tilde{\epsilon}$ terms by about 10%. This has a corresponding impact on the Gronau-London-Wyler [16, 17] and Atwood-Dunietz-Soni [14] methods to get at the CKM phase γ with the $B^+ \to \{D^0, \overline{D^0}\}K^+$ decays [9].

The decay rates for chains like the one in Fig. 2 are easy to calculate. Using Eq. 32 we find,

$$\begin{aligned} \Gamma\left[B^+ \overset{t_B}{\to} D \overset{t_D}{\to} f\right] \ = \ & e^{-\Gamma_{B^+} t_B} \left|A_{B^+\to D^0} A_{D^0\to f} + A_{B^+\to \overline{D^0}} A_{\overline{D^0}\to f}\right|^2 \times \\ & \left[\left|g_+^D\right|^2 + 2\mathrm{Re}\left\{\chi_1 g_+^{D*} g_-^D\right\} + \left|\chi_1\right|^2 \left|g_-^D\right|^2\right] \ , \end{aligned} \tag{37}$$

where

$$\chi_1 \equiv \frac{\lambda_{D\to f} + \xi_{B^+\to D}}{1 + \lambda_{D\to f}\, \xi_{B^+\to D}}. \tag{38}$$

Taking $y_D = 0$ and the small x_D approximation, the term dependent on $g_+^{D*} g_-^D$ is proportional to $\mathrm{Im}\chi_1\, x_D$. It contributes to the rate due to two independent interference effects: one related to $\mathrm{Im}\lambda_{D\to f}$, the other related to $\mathrm{Im}\xi_{B^+\to D}$ (see Eq. 41 bellow). This is easiest to understand with the aid of the parametrizations

$$\begin{aligned} A_{D^0\to K^-\pi^+} &= A \ , \qquad & A_{\overline{D^0}\to K^-\pi^+} &= -\epsilon\, A\, e^{i\Delta_D}, \\ A_{B^+\to \overline{D^0} K^+} &= B \ , \qquad & A_{B^+\to D^0 K^+} &= \tilde{\epsilon}\, e^{i\gamma}\, B\, e^{i\Delta_B}, \end{aligned} \tag{39}$$

where A, and B contain form factors, strong phases and magnitudes of CKM matrix elements. Δ_D and Δ_B are the differences of strong phases that exist between the amplitudes involving D^0 and the corresponding amplitudes involving $\overline{D^0}$. In addition, we take $q_D/p_D = e^{2i\theta_D}$, thus allowing for the fact that the new physics effects might also bring with them new phases into the $D^0 - \overline{D^0}$ mixing. Therefore,

$$\lambda_{D\to K^-\pi^+} = -\epsilon\, e^{i(2\theta_D + \Delta_D)} \quad \text{and} \quad \xi_{B^+\to D K^+} = \frac{1}{\tilde{\epsilon}}\, e^{-i(\gamma + 2\theta_D + \Delta_B)} \ . \tag{40}$$



Figure 3: The $B^+ \to DK^+ \to (X^- l^+ \nu_l)_D K^+$ cascade decay.

The term linear in x_D becomes proportional to

$$\begin{aligned}\mathrm{Im}\,\chi_1 &\propto \mathrm{Im}\,\lambda_{D\to K^-\pi^+}\left(1-|\xi_{B^+\to DK^+}|^2\right)+\mathrm{Im}\,\xi_{B^+\to DK^+}\left(1-|\lambda_{D\to K^-\pi^+}|^2\right)\\ &\sim \frac{1}{\tilde{\epsilon}^2}\left[\epsilon\sin\left(2\theta_D+\Delta_D\right)-\tilde{\epsilon}\sin\left(\gamma+2\theta_D+\Delta_B\right)\right]\ , \end{aligned} \tag{41}$$

where we have neglected higher order terms in ϵ and $\tilde{\epsilon}$. On the one hand, there is a contribution proportional to ϵx_D which arises from the $\mathrm{Im}\lambda_{D\to f}$ piece in χ_1. This involves θ_D, which vanishes in the SM, and the difference of strong phases Δ_D, which is difficult to estimate but could also be small. Obviously, these two effects are the same that appear in the direct decay $D \to K^-\pi^+$; they were studied in this context by [18], and [19, 20], respectively. On the other hand, there is a contribution proportional to $\tilde{\epsilon} x_D$ which arises from the $\mathrm{Im}\xi_{B^+\to D}$ piece in χ_1. This is *guaranteed* large even in the SM, for it involves the large CKM phase γ. Therefore, these decays may be used to probe values of $x_D \sim 10^{-2}$. This case is discussed in detail in reference [9].

Decays like $B^+ \to \{D^0, \overline{D^0}\}\pi^+$ and $B^+ \to \{D^0, \overline{D^0}\}\rho^+$ might also be useful in the search for x_D. These decay chains have much larger branching ratios than $B^+ \to \{D^0, \overline{D^0}\}K^+$. Indeed, $\mathrm{BR}[B^+ \to \overline{D^0}\rho^+] = (1.34 \pm 0.18) \times 10^{-2}$, $\mathrm{BR}[B^+ \to \overline{D^0}\pi^+] = (5.3 \pm 0.5) \times 10^{-3}$ [21], while $\mathrm{BR}[B^+ \to \overline{D^0}K^+]/\mathrm{BR}[B^+ \to \overline{D^0}\pi^+] = 0.055 \pm 0.015$ [22]. This means that the overall normalization factor $|B|^2$ in Eq. 37 is roughly 18 (46) times larger in $B^+ \to D\pi^+$ ($B^+ \to D\rho^+$) decays than it is in $B^+ \to DK^+$ decays. But, in these cases, $\tilde{\epsilon}$ is further suppressed by a factor of about 0.05 with respect to its value in the $B^+ \to \overline{D^0}K^+$ decay chains. This suppresses the $\epsilon\tilde{\epsilon}$ and $\tilde{\epsilon}x_D$ interference terms in the decay rate, meaning that these interference effects come into all the decay chains at roughly the same level. On the contrary, the term proportional to ϵx_D is the same for all the decay chains. Therefore, it should be much easier to detect in $B^+ \to D\pi^+$ and $B^+ \to D\rho^+$ decays than it is in the original $B^+ \to DK^+$ decays proposed by Meca and Silva [9]. We recall that the ϵx_D term involves $\mathrm{Im}\lambda_{D\to f}$ which also shows up in direct $D \to f$ decays. Hence, it is only detectable if the new physics also produces large CP-violating effects, θ_D, in the $D^0 - \overline{D^0}$ system [18], or if the strong phase difference, Δ_D, is considerable [19, 20].

The crucial role played by the ξ_i parameters in the $B^+ \to \{D^0, \overline{D^0}\}K^+ \to [f]_D K^+$ cascade decays is seen most vividly by taking $f = X^- l^+ \nu_l$. This case is illustrated in Fig. 3. Since the decay $\overline{D^0} \to X^- l^+ \nu_l$ is forbidden, we have $\lambda_{D\to f} = 0$, $\chi_1 = \xi_{B^+\to D}$ and, thus, the interference is determined solely by $\xi_{B^+\to D}$. Indeed, it is clear from

Fig. 3 that the only interference effect present is that arising from the clash between the $D^0-\overline{D^0}$ mixing and the decay from the B^+ *into that system*. Notice the crucial role played by the fact that there are two decay paths from B^+ into the $D^0-\overline{D^0}$ system; $B^+\to D^0$ and $B^+\to\overline{D^0}$. Both must be present in order for the decay chain to display this new type of interference.

Finally, there are $B^+\to\overline{D^0}D_s^+$ and $B^+\to\overline{D^0}D^+$ decays. However, there are no $B^+\to D^0D_s^+$ and $B^+\to D^0D^+$ decays. As a result, in Fig. 2 there is only one decay path from the initial B^+ into the intermediate state and, thus, the effects of that decay factor out in the overall decay rates. Everything works as if we had started from an initial $\overline{D^0}$ meson.

5.2 Cascade decays $B\to K$

Let us consider decay chains of the type $B_d^0\to K^0X\to[f]_KX$, where X may be J/ψ, ϕ, ω, ρ^0, π^0, etc. In this case, we have $A_{B_d^0\to\overline{K^0}}=0$ and $A_{\overline{B_d^0}\to K^0}=0$ in the SM. Therefore, the parameters needed for Eqs. 32 and 34 become

$$\begin{aligned}
N &= \left|A_{B_d^0\to K^0}A_{K^0\to f}\right|^2 \ , \\
\chi_1 &= \lambda_{K\to f} \ , \\
\chi_2 &= \frac{q_{B_d}}{p_{B_d}}\frac{p_K}{q_K}\frac{A_{\overline{B_d^0}\to\overline{K^0}}}{A_{B_d^0\to K^0}} = -\lambda_{B_d\to K_S} \ , \\
\chi_3 &= \chi_1\chi_2 \ .
\end{aligned} \tag{42}$$

We had already anticipated these results in Eq. 27, where we mentioned that the last equality on the third line holds if $|q_K/p_K|=1$. From the experimental measurement of CP violation in the $K^0-\overline{K^0}$ system, we know that this equality holds to the level of 10^{-3}.

The situation is very similar for the decay chains of the type $B_s^0\to\overline{K^0}X\to[f]_KX$. Here, $A_{B_s^0\to K^0}=0$ and $A_{\overline{B_s^0}\to\overline{K^0}}=0$ in the SM. Hence,

$$\begin{aligned}
N &= \left|A_{B_s^0\to\overline{K^0}}A_{\overline{K^0}\to f}\right|^2 \ , \\
\chi_1 &= 1/\lambda_{K\to f} \ , \\
\chi_2 &= \frac{q_{B_s}}{p_{B_s}}\frac{q_K}{p_K}\frac{A_{\overline{B_s^0}\to K^0}}{A_{B_s^0\to\overline{K^0}}} = -\lambda_{B_s\to K_S} \ , \\
\chi_3 &= \chi_1\chi_2 \ .
\end{aligned} \tag{43}$$

The articles [4, 5, 6, 7] discuss the first case, with $X=J/\psi$; while reference [8] describes the second case, also for $X=J/\psi$. As we have stressed before, in these cases, the fact that there is only one decay path from the B^0 into the $K^0-\overline{K^0}$ system implies that all the interference effects (and any CP violation therein) may be written in terms of the classic λ parameters.

To find the decay rates, we could substitute Eq. 42 or Eq. 43 into Eq. 32. However, we know that the lifetimes of the mass eigenstates of the $K^0-\overline{K^0}$ system are very different. Therefore, it is more appropriate to write our expressions in terms of γ_S and γ_L, which are the lifetimes of the short-lived (K_S) and long-lived (K_L) mass eigenstates, respectively. Since in both cases $\chi_3 = \chi_1 \chi_2$, Eq. 22 may be rewritten as

$$A\left[B^0 \overset{t_B}{\to} K \overset{t_K}{\to} f\right] \propto e^{-i\mu_H^K t_K} \left(1+\chi_1\right) \left(g_+^B + \chi_2 g_-^B\right) + e^{-i\mu_L^K t_K} \left(1-\chi_1\right) \left(g_+^B - \chi_2 g_-^B\right) . \tag{44}$$

Defining

$$e^{i\phi} \equiv - \frac{(1+\chi_1)(1-\chi_1^*)}{|(1+\chi_1)(1-\chi_1^*)|}, \tag{45}$$

the decay rate becomes

$$\begin{aligned} 4\,\Gamma\left[B^0 \overset{t_B}{\to} K \overset{t_K}{\to} f\right]/N = & \\ & e^{-\gamma_S t_K} \left|1-\chi_1\right|^2 \left|g_+^B - \chi_2 g_-^B\right|^2 + e^{-\gamma_L t_K} \left|1+\chi_1\right|^2 \left|g_+^B + \chi_2 g_-^B\right|^2 \\ & -2 e^{-\frac{\gamma_S+\gamma_L}{2} t_K} \left|1-\chi_1\right| \left|1+\chi_1\right| \left[\cos\left(\Delta m_K\, t_K - \phi\right) \left(\left|g_+^B\right|^2 - \left|\chi_2 g_-^B\right|^2\right)\right. \\ & \left. +2 \sin\left(\Delta m_K\, t_K - \phi\right) \mathrm{Im}\left(\chi_2 g_+^{B*} g_-^B\right)\right] . \end{aligned} \tag{46}$$

Now, using Eqs. 58 and recalling that χ_1 is either equal to λ_f or to $1/\lambda_f$, we notice that

$$\begin{aligned} \left|1-\chi_1\right| &\propto \left|A_{K_S \to f}\right| , \\ \left|1+\chi_1\right| &\propto \left|A_{K_L \to f}\right| . \end{aligned} \tag{47}$$

Therefore, the first term in Eq. 46 describes the decays in which the final state f has come from K_S. Taking $f = \pi^+\pi^-$, this is the dominant term for times $t_K < 1/\gamma_S$. Likewise, the second term refers to the decays in which the final state f has come from K_L. At late times, $t_K \gg 1/\gamma_S$, the cascade chain is dominated by these decays since by then the K_S component will have decayed away. The last terms describe the interference between the path going through K_S and that going through K_L. These will become important at intermediate times [4, 7].

Let us consider decays of the type $B_d \to K J/\psi$. In the SM, these decays get tree level and penguin contributions which, to an excellent approximation, share a common weak phase. Moreover, in the SM one has $\Delta\Gamma_{B_d} \sim 0$ and $|q_B/p_B| \sim 1$. As a result $|\chi_2|^2 = 1$, $\mathrm{Re}\chi_2 = -\cos 2\beta$, and $\mathrm{Im}\chi_2 = \sin 2\beta$. Also, since $\Delta\Gamma_B \sim 0$, we get from Eq. 65 that $\mathrm{Re}(g_+^{B*} g_-^B) \sim 0$. This is the origin of the standard result[4]: the pure K_S (and the pure K_L) term of Eq. 46 only measures $\sin 2\beta$, leaving a fourfold ambiguity in the determination of β. Kayser has pointed out that one can use the $K_S - K_L$ interference on the last line of Eq. 46 to get at $\cos 2\beta$, thus reducing the ambiguity [7].

---

[4] It should be stressed that the classic $B_d \to J/\psi K_S$ experiments involve, in fact, cascade decays, since the kaon will ultimately be detected through its decay into $\pi^+\pi^-$. However, since the K_L is mostly CP odd, the detected $\pi^+\pi^-$ is very unlikely to have come from a K_L ($\chi_1 \sim \eta_K = -1$ in Eqs. 13 and 46). Moreover, the experiments will select events with low t_K [23]. We see from Eq. 46 that, under these circumstances, the rate is overwhelmingly dominated by the first term, which corresponds to the pure $B_d \to J/\psi K_S$ decay [23].

One may also consider those decay chains where the primary decay is $B_d^0 \to K^0 \rho^0$, $B_d^0 \to K^0 \pi^0$, etc. If these primary decays are dominated by the top-mediated penguin diagram, then χ_2 measures again the weak phase β. However, although suppressed by $|V_{ub} V_{us}/(V_{tb} V_{ts})|$ and by color, the tree level diagram comes in with a different weak phase. The resulting hadronic uncertainties imply that these decay chains cannot compete with $B_d \to K J/\psi$ in the determination of the CKM phase β.

We now turn to decays of the type $B_s^0 \to \overline{K^0} J/\psi$. The considerations made about the $B_d \to K J/\psi$ case also apply here, with one exception. Here, the SM tree level and penguin contributions have a large weak phase difference, which turns out to be equal to β. In fact, the decay amplitudes may be parametrized as[5]

$$\begin{aligned} A_{B_s^0 \to \overline{K^0} J/\psi} &= V_{cb}^* V_{cd} T + V_{tb}^* V_{td} P \sim -\lambda^3 T + \lambda^3 e^{-i\beta} P, \\ -A_{\overline{B_s^0} \to K^0 J/\psi} &= V_{cb} V_{cd}^* T + V_{tb} V_{td}^* P \sim -\lambda^3 T + \lambda^3 e^{+i\beta} P. \end{aligned} \tag{48}$$

The symbols T and P stand for the tree level and penguin diagram contributions, with the CKM factor taken out; they include form-factors, strong phases, the gauge couplings (weak and strong, respectively), and P also includes the usual one-loop prefactor. Since the penguin diagram is not CKM suppressed with respect to the tree level diagram, $|\chi_2| = |\lambda_{B_s \to J/\psi K_S}|$ may differ from one by a few percent. Azimov and Dunietz have pointed out that one may use these decay chains to get at the elusive Δm_{B_s} mass difference [8]. It is well known that Δm_{B_s} is difficult to determine from time-dependent measurements because of the vertexing limits. These preclude measurements of x_s much larger than about 20. The situation is also bleak if we integrate the decays over t_B because the effects of x_s show up in the rate into a flavour-specific final state as $1/(1+x^2)$. This should be compared to what one gets by integrating Eq. 46 over t_B. As shown in the appendix, the result may be gotten in a straightforward way with the substitutions $|g_+^B|^2 \to G_+^B$, $|g_-^B|^2 \to G_-^B$, and $g_+^{B*} g_-^B \to G_{+-}^B$. Now, Azimov and Dunietz point out that, upon integration over t_B, the last term of Eq. 46 contains a term proportional to $\mathrm{Re}\chi_2 \, \mathrm{Im} G_{+-}^B$. And, the function $\mathrm{Im} G_{+-}^B$ shown in Eq. 67 involves $x/(1+x^2)$ rather than $1/(1+x^2)$ [8]. This increases the sensitivity to x_s considerably. Moreover, this term involves $\mathrm{Re}\chi_2$ which is nonzero even in the limit of CP conservation [8]. To appreciate this effect, we take $x_s \gg 1$, $|y_s| \ll 1$, and we assume that there is CP conservation. In that case, there will be no phase differences in Eqs. 48 and Eq. 29 reads $\chi_2 = \eta_X \eta_{B_s} \eta_K = -1$. Under these assumptions, we find

$$\begin{aligned} \frac{4\Gamma_{B_s} |q_K|^2}{|A_{B_s^0 \to \overline{K^0}}|^2} \int_0^\infty \Gamma \left[B_s^0 \xrightarrow{t_B} K \xrightarrow{t_K} f \right] dt_B = & \\ & e^{-\gamma_S t_K} \left| A_{K_S \to f} \right|^2 (1 - y_s) + e^{-\gamma_L t_K} \left| A_{K_L \to f} \right|^2 (1 + y_s) \\ & -2 e^{-\frac{\gamma_S + \gamma_L}{2} t_K} \left| A_{K_S \to f} A_{K_L \to f} \right| \sin\left(\Delta m_K \, t_K - \phi \right) \frac{1}{x_s} \ . \end{aligned} \tag{49}$$

Therefore, as pointed out by Azimov and Dunietz, the interference term is sensitive to $1/x_s$ [8].

---

[5] In these expressions, we have used the standard phase convention for the CKM matrix elements, and we have ignored the CP transformation phases of B_s and K. However, to be consistent, we have kept the $\eta_X = -1$ factor obtained in getting from the first to the second line [12].



Figure 4: The $B_d^0 \to DX \to (K^-\pi^+)_D X$ cascade decay.

We now look at the cascade decays with $B_s^0 \to \overline{K^0}\pi^0$, $B_s^0 \to \overline{K^0}\rho^0$, etc. If these decays are dominated by the tree-level diagram, then the phase of χ_2 measures the CKM phase γ. However, the penguin diagram comes in with a different weak phase and, moreover, the tree level diagram is suppressed by color and by $|V_{ub}V_{ud}/(V_{tb}V_{td})|$. The tree level and the penguin diagrams could come into χ_2 in commensurate proportions, with correspondingly large hadronic uncertainties. As a result, these decays may be of limited use. Of course, if we were to assume that there is CP conservation, then the phase of χ_2 would become well defined. But, in this case, such an assumption is not warranted for it would just hide the large theoretical errors that do exist in the determination of the phase of χ_2.

5.3 Cascade decays $B \to D$

Let us consider decays of the type $B_d^0 \to \{D^0, \overline{D^0}\}\pi^0$, $B_d \to \{D^0, \overline{D^0}\}\rho^0$, etc. Here, the decay $B_d^0 \to D^0 X$ is suppressed with respect to the decay $B_d^0 \to \overline{D^0} X$ by $\tilde{\epsilon}' = |V_{ub}V_{cd}/(V_{cb}V_{ud})|$. One might be tempted to conclude that everything follows as for the $B^+ \to \{D^0, \overline{D^0}\}X^+$ decay chains, and, in particular, that one might also use these decays to get at x_D. This is not the case, because of the $B_d^0 - \overline{B_d^0}$ mixing. In order to understand this, one should compare Fig. 2 and Fig. 4. The crucial point about Fig. 2 is that the two paths which exist in the absence of $D^0 - \overline{D^0}$ mixing are both suppressed: one by ϵ, the other by $\tilde{\epsilon}$. This is what makes the x_D corrections important; they are to be compared with ϵ and $\tilde{\epsilon}$, and not with 1 [9]. On the contrary, in the $B_d \to DX$ decays of Fig. 4 there are four paths in the absence of $D^0 - \overline{D^0}$ mixing. Two of the paths correspond to the unmixed $B_d^0 \to B_d^0$ time evolution. One path is suppressed by ϵ, the other by $\tilde{\epsilon}'$. These paths are analogous to the ones in the $B^+ \to DX^+$ decay chain. However, in Fig. 4 there are also two paths corresponding to the mixed evolution of B_d^0 into $\overline{B_d^0}$. One of these is suppressed by $\epsilon\tilde{\epsilon}'$, but the other, given schematically by

$$B_d^0 \to \overline{B_d^0} \to D^0 X \to D^0 X \to (K^-\pi^+)_D X, \tag{50}$$

is not suppressed at all. Therefore, in $B_d \to DX$ decays, the x_D corrections are to be compared with 1, not with ϵ or $\tilde{\epsilon}'$. For $x_D \sim 10^{-2}$, this correction amounts to about 1%, and may be safely neglected.

There are, however, two caveats to the analysis presented. First, if one is using the $B_d \to DX$ decays to look for precision measurements, such as effects proportional to ϵ

or $\tilde{\epsilon}'$, then one must indeed take the x_D corrections into account. Second, things would change if one could select events with $t_B \ll 1/\Delta m_{B_d}$. These times are too short for the B_d^0 to mix into $\overline{B_d^0}$ and, for them, the B_d^0 decay chain mimics that of a B^+.

We now turn to decays of the type $B_s \to DX$, where X is a meson whose quark content contains a $s\bar{s}$ piece. Examples include ϕ, η, etc. The analysis of these systems is exactly the same as for the previous case, except that the suppression of $\tilde{\epsilon}'$ is substituted here by a rather mild suppression given by $|V_{ub}V_{cs}/(V_{cb}V_{us})| = R_b$. We have used $|V_{ub}| = R_b \lambda^3$, with R_b ranging from 1/3 to 1/2 to account for the fact that $|V_{ub}|$ is in fact closer to $\lambda^3/2$ or even $\lambda^3/3$. Decays of this type have been discussed by Gronau and London [16]. As before, the presence of mixing (here the $B_s^0 - \overline{B_s^0}$ mixing) means that the cascade decays based upon $B_s \to D\phi$ are much less affected by a large value of x_D than the cascade decays started by $B^+ \to DX^+$.

5.3.1 The no-oscillation requirement

While we may obtain values of $x_D \sim 10^{-2}$ in many models of new physics, the value of x_D in the SM is much smaller than that. One might wonder what happens to the $B \to D$ cascade decays if we keep with the SM and take $x_D \sim y_D \sim 0$. In this case there is no oscillation in the $D^0 - \overline{D^0}$ system: $g_+^D = 1$ and $g_-^D = 0$. The decay amplitude in Eq. 22 becomes

$$A\left[B^0 \overset{t_B}{\to} D \overset{t_D}{\to} [f]_D\right] = A_{B^0 \to D_{\mathrm{into}\, f_D}} \left[g_+^B + \lambda_{B \to D_{\mathrm{into}\, f_D}} g_-^B\right] \ . \tag{51}$$

In writing this expression we have used Eq. 26 and noticed that Eq. 24 implies

$$A_{B^0 \to D_{\mathrm{into}\, f_D}} \equiv \langle D_{\mathrm{into}\, f_D} | B^0 \rangle = A_{B^0 \to D^0} A_{D^0 \to f_D} + A_{B^0 \to \overline{D^0}} A_{\overline{D^0} \to f_D}. \tag{52}$$

Therefore, in the absence of $D^0 - \overline{D^0}$ mixing, the decay chain $B^0 \to DX \to [f]_D X$ reduces to the direct decay of an initial B^0 into the final state $D_{\mathrm{into}\, f_D} X$. This is as expected. Indeed, we know that $D_{\mathrm{into}\, f_D}$ is the combination of D^0 and $\overline{D^0}$ that decays into the final state $[f]_D$ at time t_D. But, since there is no $D^0 - \overline{D^0}$ oscillation, that is also the state that one must overlap with the state obtained from the decay of the B meson into the $D^0 - \overline{D^0}$ system (at time $t_D = 0$).

Typically, f_D is taken to be a flavour specific final state, which picks up either D^0 or $\overline{D^0}$. In fact, we see from Eq. 24 that, in this case, $|D_{\mathrm{into}\, f_D}\rangle \propto |D^0\rangle$, or $|D_{\mathrm{into}\, f_D}\rangle \propto |\overline{D^0}\rangle$. Alternatively, f_D is taken to be a CP eigenstate, f_{cp}, with eigenvalue $e^{i\alpha_f} = \eta_f = \pm 1$. In this context, it is usually assumed that there is no CP violation in the decays of neutral D mesons into the final state f_{cp}. We can use Eqs. 16 and 24 to show that, under these conditions, $\mathcal{CP}|D_{\mathrm{into}\, f_{\mathrm{cp}}}\rangle = \eta_f |D_{\mathrm{into}\, f_{\mathrm{cp}}}\rangle$, as expected.

This no-oscillation requirement is the one behind the $B \to \{D^0, \overline{D^0}, D_{\mathrm{into}\, f_{\mathrm{cp}}}\}X$ triangle construction in the Gronau-London-Wyler method to measure the CKM phase γ [16, 17].

5.4 Cascade decays $B \to D_{\mathrm{into}\, f_D} K \to D_{\mathrm{into}\, f_D} [f]_K$.

Strictly speaking, in the $B \to DK$ decay chains we must follow three time dependences separately; those of the evolution in the $B^0 - \overline{B^0}$, $D^0 - \overline{D^0}$, and $K^0 - \overline{K^0}$ systems.

However, if we continue to assume that there is no evolution in the $D^0 - \overline{D^0}$ system, as we have done in Eq. 51, then these decay chains reduce to

$$B \to D_{\mathrm{into}\, f_D} K \to D_{\mathrm{into}\, f_D} [f_K]. \tag{53}$$

This is effectively a $B \to KX \to [f]_K X$ decay chain of the type described by Eqs. 44 and 46.

There are two cases to be considered. For $B_d \to DK$ decay chains we need the amplitude of $B_d^0 \to D^0 K^0$, which is proportional to $V_{ub}^* V_{cs}$, and the amplitude of $B_d^0 \to \overline{D^0} K^0$, which is proportional to $V_{cb}^* V_{us}$. These two amplitudes correspond to color-suppressed diagrams and differ by about R_b. They come into the amplitude of $B_d^0 \to D_{\mathrm{into}\, f_D} K$ in the proportions given by Eq. 52, which depend on the final state f_D into which the $D^0 - \overline{D^0}$ combination decays. We find,

$$\begin{aligned}
A_{B_d^0 \to D_{\mathrm{into}\, f_D} K^0} &= A_{B_d^0 \to \overline{D^0} K^0} A_{\overline{D^0} \to f_D} + A_{B_d^0 \to D^0 K^0} A_{D^0 \to f_D} \\
&\sim \lambda^3 T' A_{\overline{D^0} \to f_D} + R_b \lambda^3 e^{i\gamma} T A_{D^0 \to f_D}, \\
A_{\overline{B_d^0} \to D_{\mathrm{into}\, f_D} \overline{K^0}} &= A_{\overline{B_d^0} \to \overline{D^0}\, \overline{K^0}} A_{\overline{D^0} \to f_D} + A_{\overline{B_d^0} \to D^0 \overline{K^0}} A_{D^0 \to f_D} \\
&\sim R_b \lambda^3 e^{-i\gamma} T A_{\overline{D^0} \to f_D} + \lambda^3 T' A_{D^0 \to f_D},
\end{aligned} \tag{54}$$

where T and T' contain the relevant form factors and strong phases. Gronau and London [16] have noticed that the two amplitudes involved in these decays have roughly the same magnitude. Using this fact they proposed a triangle construction based on the comparison between $|A_{B_d^0 \to \overline{D^0} K^0}|$, $|A_{B_d^0 \to D^0 K^0}|$, $|A_{B_d^0 \to D_{\mathrm{into}\, f_{\mathrm{cp}}} K^0}|$, and the magnitudes of the CP conjugated decays. This allows the extraction of the CKM phase γ, up to discrete ambiguities. Gronau and London [16] also noted that the interference CP-violating terms in these decays allow the extraction of the unusual combination of CKM phases $2\beta + \gamma$.

The two amplitudes involved in $B_s \to DK$ decays are those of $B_s^0 \to D^0 \overline{K^0}$, proportional to $V_{ub}^* V_{cd}$, and $B_s^0 \to \overline{D^0}\, \overline{K^0}$, proportional to $V_{cb}^* V_{ud}$. The former is CKM suppressed with respect to the latter by roughly a factor of a few times 10^{-2}. This implies that triangles similar to those in the Gronau–London construction [16] would be extremely narrow, thus precluding a good measurement of γ in this case. In fact, using Eq. 52 we get

$$\begin{aligned}
A_{B_s^0 \to D_{\mathrm{into}\, f_D} \overline{K^0}} &= A_{B_s^0 \to \overline{D^0}\, \overline{K^0}} A_{\overline{D^0} \to f_D} + A_{B_s^0 \to D^0 \overline{K^0}} A_{D^0 \to f_D} \\
&\sim \lambda^2 T' A_{\overline{D^0} \to f_D} - R_b \lambda^4 e^{i\gamma} T A_{D^0 \to f_D}, \\
A_{\overline{B_s^0} \to D_{\mathrm{into}\, f_D} K^0} &= A_{\overline{B_s^0} \to \overline{D^0} K^0} A_{\overline{D^0} \to f_D} + A_{\overline{B_s^0} \to D^0 K^0} A_{D^0 \to f_D} \\
&\sim -R_b \lambda^4 e^{-i\gamma} T A_{\overline{D^0} \to f_D} + \lambda^2 T' A_{D^0 \to f_D}.
\end{aligned} \tag{55}$$

However, this strong hierarchy has its advantages too. Let us consider $f_D = f_{\mathrm{cp}}$ and assume that there is no CP violation in the decays of neutral D mesons into the final state f_{cp}. In this case Eq. 20 holds: $|A_{M^0 \to f}| = |A_{\overline{M^0} \to f}|$. By comparing Eqs. 55 with Eqs. 48 we conclude that the proposal to measure Δm_{B_s} put forth by Azimov and Dunietz [8], can be replicated by using the cascade decay $B_s \to D_{\mathrm{into}\, f_D} K \to D_{\mathrm{into}\, f_D} [f_K]$



Figure 5: The cascade decay $D^0 \to K\pi^0 \to [X^- l^+ \nu_l]_K \pi^0$.

instead. We recall that Eqs. 48 imply that $|\chi_2|$ may differ from 1 by about $|P/T|$. Here this problem is mitigated by the fact that the second amplitude in the first Eq. 55 is suppressed by $\sim 10^{-2}$.

5.5 Cascade decays $D \to K$

Let us now consider decays of the type $D^0 \to \{K^0, \overline{K^0}\}X \to [f]_K X$. These decays do not look promising. We discuss them briefly, for the sake of completeness. The decays involve the CKM matrix elements of the first two families. In the standard phase convention for the CKM matrix, these matrix elements are very approximately real. Moreover, since $\delta_K \equiv |p_K|^2 - |q_K|^2$ is known experimentally to be very small, the only rephasing-invariant quantities in $D^0 \to \{K^0, \overline{K^0}\}X \to [f]_K X$ decays which have non-negligible weak phases are those involving $q_D/p_D = e^{2i\theta_D}$, and this only if the new physics effects turn out to make θ_D large. Since the large weak phases (β and γ) are not involved, and since the kaon system is well tested experimentally, these decays can only be used to obtain the mixing parameters in the $D^0 - \overline{D^0}$ system.

However, due to the fact that the mixing of the D meson is very small (we continue to take $x_D \sim 10^{-2}$ as our reference value) this mixing can only be relevant if the amplitudes of the other paths available to get to the final state are suppressed in some way (as they are in the decay chain shown in Figs. 2 and 3). This has two important consequences. First, one of the amplitudes, $D^0 \to K^0 X$ or $D^0 \to \overline{K^0}X$, must be suppressed relatively to the other. The decays $D^0 \to K^0\pi^0$ and $D^0 \to \overline{K^0}\pi^0$ are of this kind, since

$$|A_{D^0 \to K^0\pi^0}| \sim \lambda^2 |A_{D^0 \to \overline{K^0}\pi^0}|. \tag{56}$$

Second, the final state $[f]_K$ cannot be a CP eigenstate, since one of the $K \to [f]_K$ decay amplitudes must also be suppressed relatively to the other (as in Fig. 2). This second requirement is clearly satisfied by the semileptonic decays of the neutral kaons, shown in Fig. 5. In that case $A_{\overline{K^0} \to f} = 0$.

However, the large mixing in the $K^0 - \overline{K^0}$ system will make it possible, even then, to get to the final state $(X^- l^+ \nu_l)$ with no suppression at all. This effect is similar to the one described in section 5.3 for the mixing of the B_d. So, to measure the mixing parameters in the $D^0 - \overline{D^0}$ system with this kind of decay, we will also need to select events with very small t_K. The idea is that, for very short times t_K the neutral kaons have not yet mixed appreciably, and one might probe effects which are due mainly to the mixing in the $D^0 - \overline{D^0}$ system. But, by looking at Fig. 5 we recognize that this requires a time cut of order $t_K \sim x_D/\Delta m_K$ or $t_K \sim x_D/|\Delta\Gamma_K|$ for the x_D effect to be dominant. Experimentally, $|\Delta\Gamma_K| \sim \gamma_S \sim 2\Delta m_K$. Therefore, we would need to look at times of order $t_K \sim x_D \tau_S$, where $\tau_S \sim 10^{-11}s$ is the lifetime of K_S!

This result can be checked directly in the decay rate. Eq. 57 shows the expression for the decay rate, integrated over t_D and expanded only up to the terms linear in $x_K\tau_K$ and $y_K\tau_K$. Using Eq. 32, we find[6]

$$\begin{aligned} e^{\tau_K} \int_0^\infty \Gamma\left[D^0 \overset{t_D}{\rightarrow} K \overset{t_K}{\rightarrow} f\right] dt_D \;&\propto\; 1 + \mathrm{Im}\chi_3 x_D \\ &+\; y_K\tau_K \left[\mathrm{Im}(\chi_1{\chi_3}^*)\frac{x_D}{2} - \mathrm{Im}\chi_2\frac{x_D}{2} - \mathrm{Re}\chi_1\right] \\ &+\; x_K\tau_K \left[\mathrm{Im}\chi_1 + \mathrm{Re}(\chi_1{\chi_3}^*)\frac{x_D}{2} - \mathrm{Re}\chi_2\frac{x_D}{2}\right] \end{aligned} \qquad (57)$$

We have expanded in x_D and taken $y_D = 0$. If we take the (mathematical) limit $\tau_K = 0$, then we get a decay of the type already studied by Wolfenstein [18], with a term proportional to $\mathrm{Im}\chi_3 x_D$. However, by developing Eq. 57 up to order τ_K^2 and taking $\chi_3 = \chi_1\chi_2$, we find that the terms proportional to x_D are only dominant up to times of order $t_K \sim x_D\,\mathrm{Re}\chi_2/\Delta m_K$, or $t_K \sim x_D\,\mathrm{Im}\chi_2/|\Delta\Gamma_K|$. (Notice that $\mathrm{Im}\chi_2$ can only be different from zero if the $D^0 - \overline{D^0}$ mixing phase is also different from zero [18], or if there is some strong phase involved [19, 20].) This confirms the result that we had guessed just by looking at Fig. 5. We conclude that, for times t_K larger than about $x_D\tau_S$, the effect of x_D will be at most of order 1%.

6 Conclusions

We have developed the formalism needed to study the most general cascade decays of the type $P \to M + \cdots \to f + \cdots$, which involve two neutral meson systems, P and M, in succession. The resulting decay rates exhibit the usual sources of CP violation: CPV in the mixing of neutral meson systems, probed by $|q/p| - 1$; CPV present directly in the decays, detected by $|A_{i\to f}| - |A_{\bar{i}\to\bar{f}}|$; and the CPV in the interference between the mixing in the initial neutral meson system, i, and the decay *from that system* into the final state f, measured by $\arg\lambda_{i\to f} + \arg\lambda_{i\to\bar{f}}$ (which is proportional to $\mathrm{Im}\lambda_{i\to f}$ when f is a CP eigenstate).

But, when both the $P \to M + \cdots$ and $P \to \overline{M} + \cdots$ decays are allowed, we find a new class of rephasing-invariant parameters, ξ_i, that measure the interference between the mixing in the $M^0 - \overline{M^0}$ system and the decay from the initial state (P^0 and $\overline{P^0}$) *into that system.*

We have applied this formalism to a variety of cases. The main results are the following. The proposal by Meca and Silva [9] to detect new physics in x_D through the decays $B^+ \to \{D^0, \overline{D^0}\}K^+ \to [f]_D K^+$ may be extended to the $B^+ \to \{D^0, \overline{D^0}\}\pi^+$

---

[6] It is instructive to derive this formula in a different way. Neglecting the weak phases of the first two families, $\chi_3 = \chi_1\chi_2$ and the $D^0 \to \{K^0, \overline{K^0}\}X \to [f]_K X$ decay rate may be written in a form similar to that of the $B \to K$ decay rate shown in Eq. 46. It is straightforward to use this to rederive Eq. 57.

and $B^+ \to \{D^0, \overline{D^0}\}\rho^+$ decays, which have much larger branching ratios. As we saw in Eqs. 41, there are two interference effects that probe x_D. If we neglect the strong phase differences, one is proportional to a possible CP-violating phase in the $D^0 - \overline{D^0}$ mixing due to new physics. This effect might be easier to detect with the new decays proposed here than it was in the original Meca and Silva proposal. The other interference effect exists even in the absence of new CP-violating phases in $D^0 - \overline{D^0}$ mixing, and it involves the CKM phase γ. When the branching ratios and suppression factors are taken into account, this effect comes into the new decays at roughly the same level as it does in $B^+ \to \{D^0, \overline{D^0}\}K^+$ decays.

Next, we showed that the mixing in the $B_d^0 - \overline{B_d^0}$ system implies that the same x_D effects are much less important in the $B_d \to \{D^0, \overline{D^0}\}\pi^0$ and $B_d \to \{D^0, \overline{D^0}\}\rho^0$ (unless one could apply a very stringent time cut on the B_d decays). The same applies to $B_d \to \{D^0, \overline{D^0}\}\phi$ decays. Similarly, $D \to \{K^0, \overline{K^0}\}\pi^0$ decays cannot be used to get a handle on x_D unless one were able to perform a very stringent time cut on t_K.

On the other hand, if we neglect the $D^0 - \overline{D^0}$ mixing, any $B \to DK \to [f]_D[f]_K$ decay chain (which would normally depend on three time variables, t_B, t_D, and t_K) may be analyzed as a decay chain of the type $B \to D_{\text{into } f_D} K \to D_{\text{into } f_D}[f]_K$. These chains can also be studied with the formalism developed in this article. The most interesting case is $B_s \to D_{\text{into } f_{\text{cp}}} K$. In principle, this could be used to get at Δm_{B_s} in the same way as proposed by Azimov and Dunietz with the $B_s \to J/\psi K \to J/\psi [f]_K$ decay chains.

We thank C. Meca for discussions on related subjects, and L. Lavoura for this and for reading and criticizing the manuscript. We are indebted to B. Kayser for the seminars on cascade decays that he gave in Lisbon, and which have inspired us to study this subject. This work was partially supported by the Portuguese FCT under contract Praxis/2/2.1/Fis/223/94 and by the grant CERN/P/FAE/1141/97. The work of J. P. S. is partially supported by the grant CERN/S/FAE/1164/97. The work of M. G. Santos is partially supported by the "Fundação da Universidade de Lisboa", through CFNUL.

A Time dependent functions involved in the decays of neutral meson systems

Assuming CPT invariance, the mass eigenstates of the $M^0 - \overline{M^0}$ system are related to the flavour eigenstates by

$$\begin{aligned} |M_H\rangle &= p_M|M^0\rangle + q_M|\overline{M^0}\rangle \\ |M_L\rangle &= p_M|M^0\rangle - q_M|\overline{M^0}\rangle \end{aligned} \tag{58}$$

with $|p_M|^2 + |q_M|^2 = 1$ and

$$\frac{q_M}{p_M} = \frac{\Delta m - \frac{i}{2}\Delta\Gamma}{2R_{12}} = \sqrt{\frac{R_{21}}{R_{12}}} \tag{59}$$

where $\Delta m = m_H - m_L$ (H-heavy, L-light) is positive by definition, $\Delta\Gamma = \Gamma_H - \Gamma_L$, and R_{12} is the off-diagonal matrix element in the effective time evolution in the $M^0 - \overline{M^0}$ space.

Consider a M^0 ($\overline{M^0}$) meson which is created at time $t_M = 0$ and denote by $M^0(t_M)$ ($\overline{M^0}(t_M)$) the state that it evolves into after a time t_M, measured in the rest frame of the meson M. We find

$$\begin{aligned} |M^0(t_M)\rangle &= g_+(t_M)|M^0\rangle + \frac{q_M}{p_M} g_-(t_M)|\overline{M^0}\rangle, \\ |\overline{M^0}(t_M)\rangle &= \frac{p_M}{q_M} g_-(t_M)|M^0\rangle + g_+(t_M)|\overline{M^0}\rangle, \end{aligned} \tag{60}$$

where

$$g_\pm(t) \equiv \frac{1}{2}\left(e^{-i\mu_H t} \pm e^{-i\mu_L t}\right), \tag{61}$$

$\mu_H \equiv m_H - i\Gamma_H/2$, and $\mu_L \equiv m_L - i\Gamma_L/2$. Similar expressions hold for the $P^0 - \overline{P^0}$ system.

The following formulas are useful:

$$\begin{aligned} |g_\pm(t)|^2 &= \frac{1}{4}\left[e^{-\Gamma_H t} + e^{-\Gamma_L t} \pm 2e^{-\Gamma t}\cos(\Delta m\, t)\right] && (62) \\ &= \frac{e^{-\Gamma t}}{2}\left[\cosh\frac{\Delta\Gamma t}{2} \pm \cos(\Delta m t)\right], && (63) \\ g_+^*(t)g_-(t) &= \frac{1}{4}\left[e^{-\Gamma_H t} - e^{-\Gamma_L t} - 2ie^{-\Gamma t}\sin(\Delta m\, t)\right] && (64) \\ &= -\frac{e^{-\Gamma t}}{2}\left[\sinh\frac{\Delta\Gamma t}{2} + i\sin(\Delta m t)\right]. && (65) \end{aligned}$$

Integrating over time, we obtain

$$\begin{aligned} G_\pm &\equiv \int_0^{+\infty} |g_\pm(t)|^2 dt = \frac{1}{2\Gamma}\left(\frac{1}{1-y^2} \pm \frac{1}{1+x^2}\right), && (66) \\ G_{+-} &\equiv \int_0^{+\infty} g_+^*(t)g_-(t) dt = \frac{1}{2\Gamma}\left(\frac{-y}{1-y^2} + \frac{-ix}{1+x^2}\right), && (67) \end{aligned}$$

where $x \equiv \Delta m/\Gamma$ and $y = \Delta\Gamma/(2\Gamma)$.

References

[1] J. H. Christenson, J. W. Cronin, V. L. Fitch and R. Turlay, Phys. Rev. Lett. **13**, 138 (1964).

[2] N. Cabibbo, Phys. Rev. Lett. **10**, 531 (1963); M. Kobayashi and T. Maskawa, Prog. Theor. Phys. **49**, 652 (1973).

[3] A. B. Carter and A. I. Sanda, Phys. Rev. Lett. **45**, 952 (1980); *ibid* Phys. Rev. D **23**, 1567 (1981); I. I. Bigi and A. I. Sanda, Nucl. Phys. **B193**, 85 (1981); *ibid* **B281**, 41 (1987).

[4] Ya. I. Azimov, JETP Letters **50**, 447 (1989); Phys. Rev. D **42**, 3705 (1990).

[5] G. V. Dass and K. V. L. Sarma, Int. J. Mod. Phys. **A7**, 6081 (1992); (E) *ibid* **A8**, 1183 (1993).

[6] B. Kayser and L. Stodolsky, Max-Planck Institute preprint number MPI-PHT-96-112, in hep-ph/9610522.

[7] B. Kayser, published in the proceedings *Les Arcs 1997, Electroweak interactions and unified theories,* pages 389-398. Also in hep-ph/9709382.

[8] Ya. I. Azimov and I. Dunietz, Phys. Lett. B **395**, 334 (1997).

[9] C. C. Meca and J. P. Silva, CFNUL report number CFNUL/98-03, in hep-ph/9807320. To appear in Phys. Rev. Lett. (1998).

[10] G. C. Branco, W. Grimus and L. Lavoura, Phys. Lett. B **372**, 311 (1996).

[11] I. Dunietz, Phys. Rev. D **52**, 3048 (1995).

[12] We thank Ya. I. Azimov and I. Dunietz for stressing the importance of this point, in connection with their article in [8].

[13] Z.-Z. Xing, Phys. Lett. B **371**, 310 (1996).

[14] D. Atwood, I. Dunietz, and A. Soni, Phys. Rev. Lett. **78**, 3257 (1997).

[15] For a review see, for example, Y. Nir, Nuovo Cim. **109A**, 991 (1996). Generically, these models keep $y_D \equiv \Delta\Gamma_D/(2\Gamma_D)$ exceedingly small, and, thus, we will drop it from our analysis.

[16] M. Gronau and D. London, Phys. Lett. B **253**, 483 (1991).

[17] M. Gronau and D. Wyler, Phys. Lett. B **265**, 172 (1991).

[18] L. Wolfenstein, Phys. Rev. Lett. **75**, 2460 (1995). See also T. Liu, Harvard University preprint number HUTP-94/E021, in *Proceedings of the Charm 2000 Workshop*, FERMILAB-Conf-94/190, page 375, 1994.

[19] G. Blaylock, A. Seiden, and Y. Nir, Phys. Lett. B **355**, 555 (1995).

[20] T. E. Browder and S. Pakvasa, Phys. Lett. B **383**, 475 (1996).

[21] C. Caso *et al.*, European Physical Journal **C3**, 1 (1998), and also the URL: http://pdg.lbl.gov.

[22] M. Athanas *et al.*, CLEO Collaboration, CLEO preprint number CLEO 98-2.

[23] B. Kayser, private communication.